

On the fermionic T-duality of the $AdS_4 \times \mathbb{CP}^3$ sigma-model

Ido Adam

Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut)
Am Mühlenberg 1, D-14476 Golm, Germany
idoadam@aei.mpg.de

Amit Dekel and Yaron Oz

Raymond and Beverly Sackler School of Physics and Astronomy
Tel-Aviv University, Ramat-Aviv 69978, Israel
amitde@post.tau.ac.il, yaronoz@post.tau.ac.il

ABSTRACT: In this note we consider a fermionic T-duality of the coset realization of the type IIA sigma-model on $AdS_4 \times \mathbb{CP}^3$ with respect to the three flat directions in AdS_4 , six of the fermionic coordinates and three of the $\mathbb{CP}^3$ directions. We show that the Buscher procedure fails as it leads to a singular transformation and discuss the result and its implications.

KEYWORDS: Duality in Gauge Field Theories, String Duality.

Contents

1. Introduction and summary	1
2. T-dualizing $AdS_4 \times \mathbb{CP}^3$	2
3. Discussion	5
A. The $\mathfrak{osp}(6 4)$ superalgebra	6

1. Introduction and summary

Since the $\mathcal{N} = 6$ superconformal Chern-Simons theory with matter was proposed by ABJM [1] as a dual to M-theory on $AdS_4 \times S^7/\mathbb{Z}_k$, which reduces in a certain limit to the type IIA superstring on $AdS_4 \times \mathbb{CP}^3$, much work has been devoted to understanding the properties of the ABJM field theory.

Several tree-level scattering amplitudes of the ABJM theory were computed [2] and were shown to possess a Yangian symmetry, which includes the non-local charges and the dual superconformal symmetry [3]. Some light-like polygonal Wilson loops in the ABJM theory were computed in [4] and hinted that the ABJM theory may have a scattering amplitudes/Wilson loop duality, which would further support the case in favor of the existence of dual superconformal symmetry. Additionally, a contour integral reproducing the known tree-level amplitudes has been recently proposed and was shown to have a Yangian symmetry [5]. Furthermore, a differential representation of a dual superconformal symmetry at tree-level has been constructed [6]. This representation involves variables dual to the ones parameterizing part of the R-symmetry in addition to the ones dual to the bosonic and fermionic momenta.

The corresponding findings in $\mathcal{N} = 4$ SYM in four dimensions were explained from the point of view of string theory on $AdS_5 \times S^5$ by a combination of bosonic and fermionic T-dualities, which is exact at the string tree-level [7, 8] (see [9] for a short review). Hence, it is interesting to see whether that is also the case for type IIA strings on $AdS_4 \times \mathbb{CP}^3$. Previously, it was found that the sigma-model for $AdS_4 \times \mathbb{CP}^3$, realized as the coset $\text{OSp}(6|4)/(\text{SO}(2,1) \times \text{U}(3))$ constructed in [10, 11], was not self-dual under T-duality involving both three directions in AdS_4 and six fermionic coordinates [12, 13]. In fact, one could not perform a fermionic T-duality in six fermionic isometries which together with the dualized bosonic ones form an Abelian subgroup of the whole isometry group.

In this note, in light of a suggestion that T-dualizing three isometries of $\mathbb{CP}^3$ is also required [3] and the new evidence [5, 6] from the field theory, we consider the fermionic T-duality along the three flat AdS_4 coordinates, three complex Killing vectors in $\mathbb{CP}^3$ (each one of real dimension one) as well as six of the fermionic coordinates, whose corresponding tangent-space vectors generate an Abelian subgroup of the isometry group. We show that as in the case of dualizing just in AdS_4 and the fermions, the Buscher procedure fails as it leads to a singular transformation [12].

The outline of this note is as follows: in Section 2 we apply the Buscher procedure for T-duality to the $OSp(6|4)/(SO(2,1) \times U(3))$ Green-Schwarz sigma-model describing type IIA strings on $AdS_4 \times \mathbb{CP}^3$ in a certain partial gauge-fixing and show that it fails. In Section 3 we discuss the implications of the result. The $osp(6|4)$ algebra is given in Appendix A.

2. T-dualizing $AdS_4 \times \mathbb{CP}^3$

We attempt to T-dualize $AdS_4 \times \mathbb{CP}^3$ along the directions corresponding to P_a , $Q_{l\alpha}$, R_{kl} , which form an Abelian subalgebra of the isometry group.

We assume that κ -symmetry can be partially gauge-fixed to set the six coordinates corresponding to $\hat{S}_\alpha^l$ to zero and choose the coset representative

$$g = e^{x^a P_a + \theta^{l\alpha} Q_{l\alpha} + y^{kl} R_{kl}} e^B, \quad e^B = e^{\hat{\theta}_l^\alpha \hat{Q}_\alpha^l + \xi^{l\alpha} S_{l\alpha}} y^D e^{\hat{y}_{kl} \hat{R}^{kl}}, \quad (2.1)$$

where the indices $a = 0, 1, 2$ run over the flat directions of AdS_4 , $\alpha = 1, 2$ are AdS_4 spinor indices and $l = 1, 2, 3$ are $U(3)$ fundamental representation indices (see Appendix A for further details). The Maurer-Cartan one-form is

$$K = J + j, \quad J = e^{-B}(dx^a P_a + d\theta^{l\alpha} Q_{l\alpha} + dy^{kl} R_{kl})e^B, \quad j = e^{-B}de^B. \quad (2.2)$$

Examining the algebra, one finds that the current J takes values in the space spanned by $\{P_a, Q_{l\alpha}, R_{kl}, \hat{Q}_\alpha^l, \lambda_k^l, \hat{R}^{kl}\}$, while j is valued in $\text{span}\{\hat{Q}_\alpha^l, S_{l\alpha}, \hat{S}_\alpha^l, D, M_{ab}, \lambda_k^l, \hat{R}^{kl}\}$.

Denoting the decomposition of K into the $\mathbb{Z}_4$ -invariant subspaces by $K_i \in \mathcal{H}_i$, the Green-Schwarz action takes the form

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left\{ -\frac{1}{2} \eta_{ab} J_{P_a} \bar{J}_{P_b} - j_D \bar{j}_D - 2J_{R_{kl}} (\bar{J}_{\hat{R}^{kl}} + \bar{j}_{\hat{R}^{kl}}) - 2\bar{J}_{R_{kl}} (J_{\hat{R}^{kl}} + j_{\hat{R}^{kl}}) - \right. \\ \left. - \frac{i}{2} C_{\alpha\beta} \left[J_{Q_{l\alpha}} (\bar{J}_{\hat{Q}_\beta^l} + \bar{j}_{\hat{Q}_\beta^l}) - (J_{\hat{Q}_\alpha^l} + j_{\hat{Q}_\alpha^l}) \bar{J}_{Q_{l\beta}} - j_{S_{l\alpha}} \bar{j}_{\hat{S}_\beta^l} + j_{\hat{S}_\alpha^l} \bar{j}_{S_{l\beta}} \right] \right\}. \quad (2.3)$$

We attempt to T-dualize the action by using the Buscher procedure [14, 15] by introducing the new fields A^a , $A^{l\alpha}$, A^{kl} , $\bar{A}^a$, $\bar{A}^{l\alpha}$ and $\bar{A}^{kl}$ such that the current now reads

$$J = e^{-B}(A^a P_a + A^{l\alpha} Q_{l\alpha} + A^{kl} R_{kl})e^B, \quad (2.4)$$

while j , which does not contain x^a , $\theta^{l\alpha}$ and y^{kl} , remains unmodified. In addition, the following Lagrange multiplier terms are added to the action:

$$S_L = \frac{R^2}{4\pi\alpha'} \int d^2z \left[\tilde{x}_a (\bar{\partial} A^a - \partial \bar{A}^a) + \tilde{\theta}_{l\alpha} (\bar{\partial} A^{l\alpha} - \partial \bar{A}^{l\alpha}) + \tilde{y}_{kl} (\bar{\partial} A^{kl} - \partial \bar{A}^{kl}) \right] , \quad (2.5)$$

where $\tilde{x}_a$, $\tilde{\theta}_{l\alpha}$ and $\tilde{y}_{kl}$ are Lagrange multipliers.

The T-duality is performed by integrating out the gauge fields, whose equations of motion are

$$\begin{aligned} 0 &= -\frac{1}{2}\eta_{bc}[e^{-B}P_a e^B]_{P_b} J_{P_c} + \frac{i}{2}C_{\alpha\beta} \left[[e^{-B}P_a e^B]_{Q_{l\alpha}} (J_{\hat{Q}_\beta^l} + j_{\hat{Q}_\beta^l}) - \right. \\ &\quad \left. - [e^{-B}P_a e^B]_{\hat{Q}_\alpha^l} J_{Q_{l\beta}} \right] - 2[e^{-B}P_a e^B]_{R_{kl}} (J_{\hat{R}^{kl}} + j_{\hat{R}^{kl}}) - 2[e^{-B}P_a e^B]_{\hat{R}^{kl}} J_{R_{kl}} + \partial \tilde{x}_a , \\ 0 &= -\frac{1}{2}\eta_{bc}[e^{-B}Q_{l\alpha} e^B]_{P_b} J_{P_c} + \frac{i}{2}C_{\beta\gamma} \left[[e^{-B}Q_{l\alpha} e^B]_{Q_{k\beta}} (J_{\hat{Q}_\gamma^k} + j_{\hat{Q}_\gamma^k}) - \right. \\ &\quad \left. - [e^{-B}Q_{l\alpha} e^B]_{\hat{Q}_\beta^k} J_{Q_{k\gamma}} \right] - 2[e^{-B}Q_{l\alpha} e^B]_{R_{pq}} (J_{\hat{R}^{pq}} + j_{\hat{R}^{pq}}) - 2[e^{-B}Q_{l\alpha} e^B]_{\hat{R}^{pq}} J_{R_{pq}} - \\ &\quad - \partial \tilde{\theta}_{l\alpha} , \\ 0 &= -\frac{1}{2}\eta_{bc}[e^{-B}R_{kl} e^B]_{P_b} J_{P_c} + \frac{i}{2}C_{\alpha\beta} \left[[e^{-B}R_{kl} e^B]_{Q_{p\alpha}} (J_{\hat{Q}_\beta^p} + j_{\hat{Q}_\beta^p}) - \right. \\ &\quad \left. - [e^{-B}R_{kl} e^B]_{\hat{Q}_\alpha^p} J_{Q_{p\beta}} \right] - 2[e^{-B}R_{kl} e^B]_{R_{pq}} (J_{\hat{R}^{pq}} + j_{\hat{R}^{pq}}) - 2[e^{-B}R_{kl} e^B]_{\hat{R}^{pq}} J_{R_{pq}} + \\ &\quad + \partial \tilde{y}_{kl} \end{aligned} \quad (2.6)$$

for the holomorphic fields and

$$\begin{aligned} 0 &= -\frac{1}{2}\eta_{bc}[e^{-B}P_a e^B]_{P_b} \bar{J}_{P_c} - \frac{i}{2}C_{\alpha\beta} \left[[e^{-B}P_a e^B]_{Q_{l\alpha}} (\bar{J}_{\hat{Q}_\beta^l} + \bar{j}_{\hat{Q}_\beta^l}) - [e^{-B}P_a e^B]_{\hat{Q}_\alpha^l} \bar{J}_{Q_{l\beta}} \right] - \\ &\quad - 2[e^{-B}P_a e^B]_{R_{kl}} (\bar{J}_{\hat{R}^{kl}} + \bar{j}_{\hat{R}^{kl}}) - 2[e^{-B}P_a e^B]_{\hat{R}^{kl}} \bar{J}_{R_{kl}} - \bar{\partial} \tilde{x}_a , \\ 0 &= -\frac{1}{2}\eta_{bc}[e^{-B}Q_{l\alpha} e^B]_{P_b} \bar{J}_{P_c} - \frac{i}{2}C_{\beta\gamma} \left[[e^{-B}Q_{l\alpha} e^B]_{Q_{k\beta}} (\bar{J}_{\hat{Q}_\gamma^k} + \bar{j}_{\hat{Q}_\gamma^k}) - \right. \\ &\quad \left. - [e^{-B}Q_{l\alpha} e^B]_{\hat{Q}_\beta^k} \bar{J}_{Q_{k\gamma}} \right] - 2[e^{-B}Q_{l\alpha} e^B]_{R_{pq}} (\bar{J}_{\hat{R}^{pq}} + \bar{j}_{\hat{R}^{pq}}) - 2[e^{-B}Q_{l\alpha} e^B]_{\hat{R}^{pq}} \bar{J}_{R_{pq}} + \\ &\quad + \bar{\partial} \tilde{\theta}_{l\alpha} , \\ 0 &= -\frac{1}{2}\eta_{bc}[e^{-B}R_{kl} e^B]_{P_b} \bar{J}_{P_c} - \frac{i}{2}C_{\alpha\beta} \left[[e^{-B}R_{kl} e^B]_{Q_{p\alpha}} (\bar{J}_{\hat{Q}_\beta^p} + \bar{j}_{\hat{Q}_\beta^p}) - \right. \\ &\quad \left. - [e^{-B}R_{kl} e^B]_{\hat{Q}_\alpha^p} \bar{J}_{Q_{p\beta}} \right] - 2[e^{-B}R_{kl} e^B]_{R_{pq}} (\bar{J}_{\hat{R}^{pq}} + \bar{j}_{\hat{R}^{pq}}) - 2[e^{-B}R_{kl} e^B]_{\hat{R}^{pq}} \bar{J}_{R_{pq}} - \\ &\quad - \bar{\partial} \tilde{y}_{kl} \end{aligned} \quad (2.7)$$

for the anti-holomorphic ones. (The complexity of the equations arises from the fact that, unlike in the $AdS_5 \times S^5$ case, J is valued in a space larger than the one that is actually dualized.)

For the purpose of solving these equations, the properties of the field-dependent group-theoretic factors must be understood. In particular, it should be checked whether the coefficients of the gauge fields have non-trivial kernels.

In order to do so, we resort to explicitly expressing the currents in terms of the coordinates. We denote $C \equiv \hat{\theta}_l^\alpha \hat{Q}_\alpha^l + \xi^{l\alpha} S_{l\alpha}$ and examine the commutators

$$\begin{aligned} [P_a, C] &= -\frac{i}{\sqrt{2}} \gamma_{a\alpha}{}^\beta \xi^{l\alpha} Q_{l\beta} \equiv \Xi_a^{Pl\beta} Q_{l\beta} , \\ [Q_{l\beta}, C] &= \frac{1}{\sqrt{2}} (\gamma^a C)_{\beta\alpha} \hat{\theta}_l^\alpha P_a + \frac{1}{\sqrt{2}} C_{\beta\alpha} \xi^{k\alpha} R_{lk} \equiv \Theta_{l\beta}^{Qa} P_a + \Xi_\beta^{Qk} R_{lk} \equiv M_{l\beta} , \\ [R_{kl}, C] &= -\frac{i}{\sqrt{2}} (\hat{\theta}_l^\alpha \delta_k^p - \hat{\theta}_k^\alpha \delta_l^p) Q_{p\alpha} \equiv \Theta_{kl}^{Rp\alpha} Q_{p\alpha} . \end{aligned} \quad (2.8)$$

We further define

$$N_{l\alpha}{}^{k\beta} = \Theta_{l\alpha}^{Qa} \Xi_a^{Pk\beta} + \Xi_\alpha^{Qp} \Theta_{pl}^{Rk\beta} \quad (2.9)$$

and note that $[M_{l\alpha}, C] = N_{l\alpha}{}^{k\beta} Q_{k\beta}$ and $[Q_{l\alpha}, C] = M_{l\alpha}$. Using the formula $e^{-B} A e^B = A + [A, B] + \frac{1}{2!} [[A, B], B] + \dots$, we get

$$\begin{aligned} e^{-C} (dx^a P_a + d\theta^{l\alpha} Q_{l\alpha} + dy^{kl} R_{kl}) e^C &= dx^a P_a + dy^{kl} R_{kl} + \\ &+ (dx^a \Xi_a^{Pl\alpha} + dy^{pq} \Theta_{pq}^{Rl\alpha}) \left[\left(\frac{\cosh \sqrt{N} - 1}{N} \right)_{l\alpha}{}^{k\beta} M_{k\beta} + \left(\frac{\sinh \sqrt{N}}{\sqrt{N}} \right)_{l\alpha}{}^{k\beta} Q_{k\beta} \right] + \\ &+ d\theta^{l\alpha} \left[\left(\frac{\sinh \sqrt{N}}{\sqrt{N}} \right)_{l\alpha}{}^{k\beta} M_{k\beta} + (\cosh \sqrt{N})_{l\alpha}{}^{k\beta} Q_{k\beta} \right] . \end{aligned} \quad (2.10)$$

Finally, conjugating with $y^D e^{\hat{y}_{kl} \hat{R}^{kl}}$ yields the current

$$\begin{aligned} J &= \frac{dx^a}{y} P_a + dy^{kl} (R_{kl} + 2i\sqrt{2} \hat{y}_{kq} \lambda_l^q + 2\hat{y}_{kq} \hat{y}_{ln} \hat{R}^{qn}) + \\ &+ \left[(dx^a \Xi_a^{Pl\alpha} + dy^{pq} \Theta_{pq}^{Rl\alpha}) \left(\frac{\cosh \sqrt{N} - 1}{N} \right)_{l\alpha}{}^{k\beta} + d\theta^{l\alpha} \left(\frac{\sinh \sqrt{N}}{\sqrt{N}} \right)_{l\alpha}{}^{k\beta} \right] \times \\ &\times \left[\tilde{M}_{k\beta} + i\sqrt{2} \Xi_\beta^{Qm} (\hat{y}_{kq} \lambda_m^q - \hat{y}_{mq} \lambda_k^q) + \Xi_\beta^{Qr} (\hat{y}_{kq} \hat{y}_{rn} - \hat{y}_{rq} \hat{y}_{kn}) \hat{R}^{qn} \right] + \\ &+ \frac{1}{y^{1/2}} \left[(dx^a \Xi_a^{Pl\alpha} + dy^{pq} \Theta_{pq}^{Rl\alpha}) \left(\frac{\sinh \sqrt{N}}{\sqrt{N}} \right)_{l\alpha}{}^{k\beta} + d\theta^{l\alpha} (\cosh \sqrt{N})_{l\alpha}{}^{k\beta} \right] \times \\ &\times (Q_{k\beta} + i\sqrt{2} \hat{y}_{pk} \hat{Q}_\beta^p) , \end{aligned} \quad (2.11)$$

where $\tilde{M}_{k\beta} \equiv y^{-D} M_{k\beta} y^D = \frac{1}{y} \Theta_{l\alpha}^{Qa} P_a + \Xi_\alpha^{Ql} R_{kl}$.

Unfortunately, j is even more complicated. However, before plunging into its computation in a closed form it is worthwhile to examine it to the lowest order in $\hat{\theta}_l^\alpha$ and $\xi^{l\alpha}$. Doing so yields,

$$j = \frac{d\hat{\theta}_l^\alpha}{y^{1/2}} \hat{Q}_\alpha^l + y^{1/2} d\xi^{l\alpha} S_{l\alpha} - i\sqrt{2} y^{1/2} \hat{y}_{kl} d\xi^{l\alpha} \hat{S}_\alpha^k + \frac{dy}{y} D + d\hat{y}_{pq} \hat{R}^{pq} + O(\hat{\theta}_l^\alpha, \xi^{l\alpha}) . \quad (2.12)$$

Having the currents, we can take a look at the action to lowest order in $\hat{\theta}_l^\alpha$ and $\xi^{l\alpha}$:

$$\begin{aligned}
S = \frac{R^2}{4\pi\alpha'} \int d^2z \Big\{ & -\frac{1}{2}\eta_{ab}\frac{\partial x^a\bar{\partial}x^b}{y^2} - \frac{\partial y\bar{\partial}y}{y^2} - 2\partial y^{kl}(2\hat{y}_{pk}\hat{y}_{ql}\bar{\partial}y^{pq} + \bar{\partial}\hat{y}_{kl}) - \\
& - 2\bar{\partial}y^{kl}(2\hat{y}_{pk}\hat{y}_{ql}\partial y^{pq} + \partial\hat{y}_{kl}) - \frac{i}{2y}C_{\alpha\beta}\Big[\partial\theta^{l\alpha}(i\sqrt{2}\hat{y}_{kl}\bar{\partial}\theta^{k\beta} + \bar{\partial}\hat{\theta}_l^\beta) - \\
& - (i\sqrt{2}\hat{y}_{kl}\partial\theta^{k\alpha} + \partial\hat{\theta}_l^\alpha)\bar{\partial}\theta^{l\beta}\Big] + \frac{i}{2}yC_{\alpha\beta}(-i\sqrt{2}\hat{y}_{lk}\partial\xi^{l\alpha}\bar{\partial}\xi^{k\beta} + i\sqrt{2}\hat{y}_{lk}\partial\xi^{k\alpha}\bar{\partial}\xi^{l\beta})\Big\}.
\end{aligned} \tag{2.13}$$

The term quadratic in the $\theta^{l\alpha}$ derivatives is multiplied by a three-dimensional antisymmetric matrix, whose rank is two, and the higher order terms in $\hat{\theta}_l^\alpha$ and $\xi^{l\alpha}$ cannot make the matrix's kernel trivial. Thus the term quadratic in the fermionic gauge fields in the dualized action will be multiplied by a singular matrix and the fermionic gauge fields will be multiplied by a singular matrix in the equations of motion — one cannot T-dualize all the six fermionic coordinates.

Since the obstruction to T-dualizing the fermionic coordinates is at the zeroth order in the spectator fermions, it appears that modifying the κ -symmetry gauge-fixing of these fermionic degrees of freedom would not change the above conclusion.

3. Discussion

We showed that the application of the Buscher T-duality procedure to the coset $\text{OSp}(6|4)/(\text{SO}(2,1) \times \text{U}(3))$ fails when dualizing along the AdS_4 flat directions, three of the (real) $\mathbb{CP}^3$ directions and six fermionic directions. There are several ways to explain this apparent tension between the field theory tree-level evidence and the sigma-model analysis.

The simplest and most obvious explanation is that the dual superconformal symmetry exists only in the weakly-coupled field theory description and breaks down at the strong-coupling regime, which is described by the string theory dual. A second possibility is that in this case the dual superconformal symmetry is not related to the ordinary superconformal symmetry by a T-duality transformation but in a more intricate way.

A third possibility is that the coset formulation does not capture the entire superstring description. The coset is obtained by a partial gauge-fixing of the κ -symmetry of the full $AdS_4 \times \mathbb{CP}^3$ sigma-model [16] by setting the fermionic coordinates corresponding to the eight broken supersymmetries to zero. However, as noted in [16], this gauge-fixing is not compatible with all the possible string configurations. Thus, it does not have a representation for certain field theory operators, which might amount to a (possibly inconsistent) truncation of the field theory that does not preserve the dual superconformal symmetry. A way to resolve this issue could be to use a better gauge-fixing of the κ -symmetry as proposed in [16, 13].

Acknowledgments

We would like to thank Y-t. Huang and A. E. Lipstein for sharing a draft of their paper [6] with us before its publication. I.A. is supported in part by the German-Israeli Project cooperation (DIP H.52) and the German-Israeli Fund (GIF).

A. The $\text{osp}(6|4)$ superalgebra

The $\text{osp}(6|4)$ algebra's commutation relations in the $\text{so}(1,2) \oplus \text{u}(3)$ basis are given by

$$[\lambda_k^l, \lambda_m^n] = \frac{i}{\sqrt{2}}(\delta_m^l \lambda_k^n - \delta_k^n \lambda_m^l), \quad (\text{A.1})$$

$$[\lambda_k^l, R_{mn}] = \frac{i}{\sqrt{2}}(\delta_m^l R_{kn} - \delta_n^l R_{km}), \quad [\lambda_l^k, \hat{R}^{pq}] = -\frac{i}{\sqrt{2}}(\delta_l^p \hat{R}^{kq} - \delta_l^q \hat{R}^{kp}) \quad (\text{A.2})$$

$$[R_{mn}, R_{kl}] = 0, \quad [R_{mn}, \hat{R}^{kl}] = \frac{i}{\sqrt{2}}(\delta_m^k \lambda_n^l - \delta_m^l \lambda_n^k - \delta_n^k \lambda_m^l + \delta_n^l \lambda_m^k) \quad (\text{A.3})$$

$$[P_a, P_b] = 0, \quad [K_a, K_b] = 0, \quad [P_a, K_b] = \eta_{ab} D - M_{ab} \quad (\text{A.4})$$

$$[M_{ab}, M_{cd}] = \eta_{ac} M_{bd} + \eta_{bd} M_{ac} - \eta_{ad} M_{bc} - \eta_{bc} M_{ad} \quad (\text{A.5})$$

$$[M_{ab}, P_c] = \eta_{ac} P_b - \eta_{bc} P_a, \quad [M_{ab}, K_c] = \eta_{ac} K_b - \eta_{bc} K_a \quad (\text{A.6})$$

$$[D, P_a] = P_a, \quad [D, K_a] = -K_a, \quad [D, M_{ab}] = 0 \quad (\text{A.7})$$

$$[D, Q_{l\alpha}] = \frac{1}{2} Q_{l\alpha}, \quad [D, S_{l\alpha}] = -\frac{1}{2} S_{l\alpha} \quad (\text{A.8})$$

$$[P_a, Q_{l\alpha}] = 0, \quad [K_a, S_{l\alpha}] = 0 \quad (\text{A.9})$$

$$[P_a, S_{l\alpha}] = -\frac{i}{\sqrt{2}}(\gamma_a)_\alpha^\beta Q_{l\beta}, \quad [K_a, Q_{l\alpha}] = \frac{i}{\sqrt{2}}(\gamma_a)_\alpha^\beta S_{l\beta} \quad (\text{A.10})$$

$$[M_{ab}, Q_{l\alpha}] = -\frac{i}{2}(\gamma_{ab})_\alpha^\beta Q_{l\beta}, \quad [M_{ab}, S_{l\alpha}] = -\frac{i}{2}(\gamma_{ab})_\alpha^\beta S_{l\beta} \quad (\text{A.11})$$

$$[R_{kl}, \hat{Q}_\alpha^p] = \frac{i}{\sqrt{2}}(\delta_l^p Q_{k\alpha} - \delta_k^p Q_{l\alpha}), \quad [R_{kl}, \hat{S}_\alpha^p] = -\frac{i}{\sqrt{2}}(\delta_l^p S_{k\alpha} - \delta_k^p S_{l\alpha}) \quad (\text{A.12})$$

$$[\hat{R}^{kl}, Q_{p\alpha}] = -\frac{i}{\sqrt{2}}(\delta_p^l \hat{Q}_\alpha^k - \delta_p^k \hat{Q}_\alpha^l), \quad [\hat{R}^{kl}, S_{p\alpha}] = \frac{i}{\sqrt{2}}(\delta_p^l \hat{S}_\alpha^k - \delta_p^k \hat{S}_\alpha^l) \quad (\text{A.13})$$

$$[\lambda_k^l, Q_{p\alpha}] = \frac{i}{\sqrt{2}}\delta_p^l Q_{k\alpha}, \quad [\lambda_k^l, S_{p\alpha}] = \frac{i}{\sqrt{2}}\delta^{pl} S_{k\alpha} \quad (\text{A.14})$$

$$[\lambda_k^l, \hat{Q}_\alpha^p] = -\frac{i}{\sqrt{2}}\delta_k^p \hat{Q}_\alpha^l, \quad [\lambda_k^l, \hat{S}_\alpha^p] = -\frac{i}{\sqrt{2}}\delta_k^p \hat{S}_\alpha^l \quad (\text{A.15})$$

$$\{Q_{l\alpha}, Q_{k\beta}\} = 0, \quad \{Q_{l\alpha}, \hat{Q}_\beta^k\} = -\frac{1}{\sqrt{2}}\delta_l^k (\gamma^a C)_{\alpha\beta} P_a \quad (\text{A.16})$$

$$\{S_{l\alpha}, S_{k\beta}\} = 0, \quad \{S_{l\alpha}, \hat{S}_\beta^k\} = -\frac{1}{\sqrt{2}}\delta_l^k (\gamma^a C)_{\alpha\beta} K_a \quad (\text{A.17})$$

$$\{Q_{l\alpha}, S_{k\beta}\} = -\frac{1}{\sqrt{2}}C_{\alpha\beta}R_{lk}, \quad \{\hat{Q}_\alpha^l, \hat{S}_\beta^k\} = -\frac{1}{\sqrt{2}}C_{\alpha\beta}\hat{R}^{lk} \quad (\text{A.18})$$

$$\{Q_{l\alpha}, \hat{S}_\beta^k\} = -i\frac{1}{2}\delta_l^k(C_{\alpha\beta}D + i\frac{1}{2}(\gamma^{ab}C)_{\alpha\beta}M_{ab}) + \frac{1}{\sqrt{2}}C_{\alpha\beta}\lambda_l^k \quad (\text{A.19})$$

$$\{\hat{Q}_\alpha^l, S_{k\beta}\} = i\frac{1}{2}\delta_k^l(C_{\alpha\beta}D - i\frac{1}{2}(\gamma^{ab}C)_{\alpha\beta}M_{ab}) + \frac{1}{\sqrt{2}}C_{\alpha\beta}\lambda_k^l \quad (\text{A.20})$$

The indices take the values $k, l = 1, \dots, 3$, the $\mathbf{3}$ of $\text{u}(3)$, $a, b = 0, 1, 2$ are the $\mathbf{3}$ of $\text{so}(1, 2)$ and $\alpha, \beta, \dots = 1, 2$ are the $\text{so}(2, 1)$ spinors, and $\eta = \text{diag}(-, +, +)$. The generators satisfy the following relations under complex conjugation $R_{kl}^* = \hat{R}^{kl}$, $\lambda_k^l = \lambda_l^{*k}$, $\hat{Q}_\alpha^l = (Q_{l\alpha})^*$ and $\hat{S}_\alpha^l = (S_{l\alpha})^*$. The $(\gamma_a)_\alpha^\beta$ are the Dirac matrices of $\text{so}(1, 2)$, and $\gamma_{ab} = \frac{i}{2}[\gamma_a, \gamma_b]$. We raise and lower spinor indices using $C_{\alpha\beta} = \epsilon_{\alpha\beta}$, $\psi_\alpha = \psi^\beta \epsilon_{\beta\alpha}$, $\psi^\alpha = \epsilon^{\alpha\beta} \psi_\beta$, where $\epsilon_{12} = -\epsilon_{21} = \epsilon^{12} = -\epsilon^{21} = 1$.

The bilinear forms are given by

$$\begin{aligned} \text{Str}(R_{kl}, \hat{R}^{pq}) &= \delta_k^q \delta_l^p - \delta_k^p \delta_l^q, \\ \text{Str}(\lambda_k^l, \lambda_p^q) &= -\delta_k^q \delta_l^p, \\ \text{Str}(Q_{l\alpha}, \hat{S}_\beta^k) &= i\delta_l^k C_{\alpha\beta}, \\ \text{Str}(S_{l\alpha}, \hat{Q}_\beta^k) &= -i\delta_k^l C_{\alpha\beta}, \\ \text{Str}(P_a, K_b) &= -\eta_{ab}, \\ \text{Str}(D, D) &= -1, \\ \text{Str}(M_{ab}, M_{cd}) &= \eta_{ac}\eta_{bd} - \eta_{ad}\eta_{bc}. \end{aligned} \quad (\text{A.21})$$

The $\mathbb{Z}_4$ subspaces with the invariant locus of $\text{U}(3) \times \text{SO}(3, 1)$ which gives the semi-symmetric space $\text{AdS}_4 \times \mathbb{CP}^3$ are

$$\begin{aligned} \mathcal{H}_0 &= \{P_a - K_a, M_{ab}, \lambda_k^l\}, \\ \mathcal{H}_1 &= \{Q_{l\alpha} - S_{l\alpha}, \hat{Q}_\alpha^l - \hat{S}_\alpha^l\}, \\ \mathcal{H}_2 &= \{P_a + K_a, D, R_{kl}, \hat{R}^{kl}\}, \\ \mathcal{H}_3 &= \{Q_{l\alpha} + S_{l\alpha}, \hat{Q}_\alpha^l + \hat{S}_\alpha^l\}. \end{aligned} \quad (\text{A.22})$$

References

- [1] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, “N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals,” *JHEP* **10** (2008) 091, [arXiv:0806.1218 \[hep-th\]](#).
- [2] A. Agarwal, N. Beisert, and T. McLoughlin, “Scattering in Mass-Deformed $N \geq 4$ Chern-Simons Models,” *JHEP* **06** (2009) 045, [arXiv:0812.3367 \[hep-th\]](#).
- [3] T. Bargheer, F. Loebbert, and C. Meneghelli, “Symmetries of Tree-level Scattering Amplitudes in N=6 Superconformal Chern-Simons Theory,” [arXiv:1003.6120 \[hep-th\]](#).
- [4] J. M. Henn, J. Plefka, and K. Wiegandt, “Light-like polygonal Wilson loops in 3d Chern-Simons and ABJM theory,” [arXiv:1004.0226 \[hep-th\]](#).

- [5] S. Lee, “Yangian Invariant Scattering Amplitudes in Super-Chern- Simons Theory,” [arXiv:1007.4772 \[hep-th\]](#).
- [6] Y.-t. Huang and A. E. Lipstein, “Dual Superconformal Symmetry of N=6 Chern-Simons Theory,” [arXiv:1008.0041 \[hep-th\]](#).
- [7] N. Berkovits and J. Maldacena, “Fermionic T-Duality, Dual Superconformal Symmetry, and the Amplitude/Wilson Loop Connection,” *JHEP* **09** (2008) 062, [arXiv:0807.3196 \[hep-th\]](#).
- [8] N. Beisert, R. Ricci, A. A. Tseytlin, and M. Wolf, “Dual Superconformal Symmetry from AdS₅ x S⁵ Superstring Integrability,” [arXiv:0807.3228 \[hep-th\]](#).
- [9] N. Beisert, “T-Duality, Dual Conformal Symmetry and Integrability for Strings on AdS₅ × S⁵,” *Fortschr. Phys.* **57** (2009) 329–337, [arXiv:0903.0609 \[hep-th\]](#).
- [10] j. Stefanski, B., “Green-Schwarz action for Type IIA strings on AdS₄ × CP³,” *Nucl. Phys.* **B808** (2009) 80–87, [arXiv:0806.4948 \[hep-th\]](#).
- [11] G. Arutyunov and S. Frolov, “Superstrings on AdS₄ × CP³ as a Coset Sigma-model,” *JHEP* **09** (2008) 129, [arXiv:0806.4940 \[hep-th\]](#).
- [12] I. Adam, A. Dekel, and Y. Oz, “On Integrable Backgrounds Self-dual under Fermionic T- duality,” *JHEP* **04** (2009) 120, [arXiv:0902.3805 \[hep-th\]](#).
- [13] P. A. Grassi, D. Sorokin, and L. Wulff, “Simplifying superstring and D-brane actions in AdS(4) x CP(3) superbackground,” *JHEP* **08** (2009) 060, [arXiv:0903.5407 \[hep-th\]](#).
- [14] T. H. Buscher, “A Symmetry of the String Background Field Equations,” *Phys. Lett.* **B194** (1987) 59.
- [15] T. H. Buscher, “Path Integral Derivation of Quantum Duality in Nonlinear Sigma Models,” *Phys. Lett.* **B201** (1988) 466.
- [16] J. Gomis, D. Sorokin, and L. Wulff, “The complete AdS(4) x CP(3) superspace for the type IIA superstring and D-branes,” [arXiv:0811.1566 \[hep-th\]](#).